

## Birds can 'read' the Earth's magnetic signature well enough to get back on course

*Date:* February 12, 2021

*Source:* Bangor University

*Summary:* Birdwatchers get excited when 'rare' migratory birds makes landfall having been blown beyond their normal range. But these are rare for a reason; most birds that have made the journey before are able to correct for large displacements and find their final destination. Now new research shows how birds displaced in this way are able to navigate back to their migratory route and gives us an insight into how they accomplish this feat.

### FULL STORY

---

Birdwatchers get very excited when a 'rare' migratory bird makes landfall having been blown off-course and flown beyond its normal range. But these are rare for a reason; most birds that have made the journey before are able to correct for large displacements and find their final destination.

Now, new research by an international team shows for the first time, how birds displaced in this way are able to navigate back to their migratory route and gives us an insight into how they accomplish this feat.

Writing in *Current Biology*, the team from Bangor and Keele Universities describe how reed warblers can navigate from a 'magnetic position' beyond what they have experienced in their normal migration route, back towards that correct route.

Different parts of the Earth have a distinct 'geomagnetic signature' according to their location. This is a combination of the strength of the geomagnetic field, the magnetic inclination or the dip angle between magnetic field lines and the horizon and the magnetic declination, or the angle between directions to the geographic and magnetic North poles.

Adult birds already familiar with their migration route, and its general magnetic signatures, were held in captivity for a short period before being released back into the wild, and exposed to a simulation of the earth's magnetic signature at a location thousands of miles beyond the birds' natural migratory corridor.

Despite remaining physically located at their capture site and experiencing all other sensory clues about their location, including starlight and the sights, smell and sounds of their actual location, the birds still showed the urge to begin their journey as though they were in the location suggested by the magnetic signal they were experiencing.

They oriented themselves to fly in a direction which would lead them 'back' to their migratory path from the location suggested to them by the magnetic signals they were experiencing.

This shows that the earth's magnetic field is the key factor in guiding reed warblers when they are blown off course.

"The overriding impulse was to respond to the magnetic information they were receiving," explained Richard Holland of Bangor University's School of Natural Sciences.

What our current work shows is that birds are able to sense that they are beyond the bounds of the magnetic fields that are familiar to them from their year-round movements, and are able to extrapolate their position sufficiently from the signals. This fascinating ability enables bird to navigate towards their normal migration route."

Dr Dmitry Kishkinev of Keele University's School of Life Sciences explained:

"What these birds are achieving is "true navigation." In other words, they are able to return to a known goal after displacement to a completely unknown location without relying on familiar surroundings, cues that emanate from the destination, or information collected during the outward journey."

Florian Packmor of Bangor University added: "We have already shown that the reed warblers use the same magnetic cues experienced within their natural range, but this study shows that they can extrapolate what they understand about how the magnetic field varies in space far beyond any previous experience they have had."

But questions remain about whether the birds have an accurate 'map' or are just using a 'rule of thumb' measurement to judge the general direction of travel needed to get back on course.

The Eurasian reed warbler was selected for the research, but the findings could probably be applied to other migrating songbirds.

---

### Story Source:

Materials provided by **Bangor University**. *Note: Content may be edited for style and length.*

---

### Journal Reference:

1. Dmitry Kishkinev et al. **Navigation by extrapolation of geomagnetic cues in a migratory songbird**. *Current Biology*, 2021 DOI: 10.1016/j.cub.2021.01.051

---

### Cite This Page:

MLA

APA

Chicago

---

Bangor University. "Birds can 'read' the Earth's magnetic signature well enough to get back on course." ScienceDaily. ScienceDaily, 12 February 2021. <[www.sciencedaily.com/releases/2021/02/210212111917.htm](http://www.sciencedaily.com/releases/2021/02/210212111917.htm)>.

---

### RELATED STORIES

#### Most Migratory Birds Rely on a Greening World

Oct. 27, 2020 — A new study confirms that most birds -- but not all -- synchronize their migratory movements with seasonal changes in vegetation greenness. This is the first study of its kind to cover the Western ...

#### Birds Migrate Away from Diseases

Apr. 10, 2018 — In a unique study, researchers have mapped the origins of migratory birds. They used the results to investigate and discover major differences in the immune systems of sedentary and migratory birds. ...

#### Even Non-Migratory Birds Use a Magnetic Compass

May 18, 2017 — Not only migratory birds use a built-in magnetic compass to navigate correctly. A new study shows that non-migratory birds also are able to use a built-in compass to orient themselves using the ...

#### Shorter Telomeres Reveal Stress in Migratory Birds

Aug. 3, 2016 — The stress of birds' continent-spanning annual migrations, it appears, leads to faster aging and a potentially earlier death. A new study reveals that telomeres, structures on the ends of chromosomes ...